

Greedoids on Vertex Sets of Unicycle Graphs

Vadim E. Levit

Department of Computer Science and Mathematics
Ariel University Center of Samaria, Israel
levitv@ariel.ac.il

Eugen Mandrescu

Department of Computer Science
Holon Institute of Technology, Israel
eugen_m@hit.ac.il

Abstract

A *maximum stable set* in a graph G is a stable set of maximum size. S is a *local maximum stable set* of G , and we write $S \in \Psi(G)$, if S is a maximum stable set of the subgraph spanned by $S \cup N(S)$, where $N(S)$ is the neighborhood of S . G is a *unicycle graph* if it owns only one cycle. In [10] we have shown that the family $\Psi(T)$ of a forest T forms a greedoid on its vertex set. Bipartite, triangle-free, and well-covered graphs G whose $\Psi(G)$ form greedoids were analyzed in [11, 12, 16], respectively.

In this paper we characterize the unicycle graphs whose families of local maximum stable sets form greedoids.

Keywords: unicycle graph, tree, bipartite graph, König-Egerváry graph, local maximum stable set, greedoid, uniquely restricted maximum matching.

1 Introduction

Throughout this paper $G = (V, E)$ is a simple (i.e., a finite, undirected, loopless and without multiple edges) graph with vertex set $V = V(G)$ and edge set $E = E(G)$. If $X \subset V$, then $G[X]$ is the subgraph of G induced by X , and by $G - W$ we mean the subgraph $G[V - W]$, where $W \subset V(G)$. The graph G is *unicycle* if it owns only one cycle. The *neighborhood* of a vertex $v \in V$ is the set $N(v) = \{w : w \in V, vw \in E\}$. If $|N(v)| = 1$, then v is a *pendant vertex*. We denote the *neighborhood* of the set $A \subset V$ by $N_G(A) = \{v \in V - A : N(v) \cap A \neq \emptyset\}$ and its *closed neighborhood* by $N_G[A] = A \cup N_G(A)$, or shortly, $N(A)$ and $N[A]$, if there is no ambiguity.

By K_n, C_n we mean the complete graph on $n \geq 1$ vertices, and the chordless cycle on $n \geq 3$ vertices, respectively.

A *stable set* in G is a set of pairwise non-adjacent vertices. A stable set of maximum size will be referred to as a *maximum stable set* of G , and the *stability number* of G , denoted by $\alpha(G)$, is the cardinality of a maximum stable set in G . By $\Omega(G)$ we denote the family of all maximum stable sets of the graph G .

A set $A \subseteq V(G)$ is a *local maximum stable set* of G if A is a maximum stable set in the subgraph induced by $N[A]$, i.e., $A \in \Omega(G[N[A]])$, [10]. Let $\Psi(G)$ stand for the family of all local maximum stable sets of G . For instance, any set S consisting of only pendant vertices belongs to $\Psi(G)$, while the converse is not generally true; e.g., the set $\{e, g\} \in \Psi(G)$ contains no pendant vertex, where G is the graph in Figure 1.




Figure 1: A graph having various local maximum stable sets.

Clearly, not any stable set of a graph G is included in some maximum stable set of G . For example, there is no $S \in \Omega(G)$ such that $\{b, d, h\} \subset S$, where G is the graph presented in Figure 1. In [18], Nemhauser and Trotter Jr. showed that every local maximum stable set of a graph can be enlarged to one of its maximum stable sets.

A *matching* in a graph $G = (V, E)$ is a set of edges $M \subseteq E$ such that no two edges of M share a common vertex. We denote the size of a *maximum matching* (a matching of maximum cardinality) by $\mu(G)$. Let us recall that G is a *König-Egerváry graph* provided $\alpha(G) + \mu(G) = |V(G)|$, [2, 19]. It is known that every bipartite graph is a König-Egerváry graph [3, 6].

A *greedoid*, [1, 5], is a pair $(V, \mathcal{F})$, where $\mathcal{F} \subseteq 2^V$ is a non-empty set system satisfying the following conditions:

accessibility: for every non-empty $X \in \mathcal{F}$, there is an $x \in X$ such that $X - \{x\} \in \mathcal{F}$;

exchange: for $X, Y \in \mathcal{F}$, $|X| = |Y| + 1$, there is an $x \in X - Y$ such that $Y \cup \{x\} \in \mathcal{F}$.

The following theorem shows that it is enough to prove that $\Psi(G)$ satisfies the accessibility property, in order to validate that $\Psi(G)$ is a greedoid.

Theorem 1.1 [17] *If the family $\Psi(G)$ satisfies the accessibility property, then it satisfies the exchange property as well.*

Clearly, $\Omega(G) \subseteq \Psi(G)$ holds for any graph G .

If $S \in \Psi(G)$, $|S| = k \geq 2$, then sometimes there exists a chain

$$\{x_1\} \subset \{x_1, x_2\} \subset \dots \subset \{x_1, \dots, x_{k-1}\} \subset \{x_1, \dots, x_{k-1}, x_k\} = S$$

such that $\{x_1, x_2, \dots, x_j\} \in \Psi(G)$, for all $j \in \{1, \dots, k-1\}$; such a chain is called an *accessibility chain* for S , [11]. It is evident that x_1 must be a simplicial vertex, i.e., a vertex whose closed neighborhood induces a complete graph in G (in particular, any pendant vertex is also simplicial).




Figure 2: G_1 and G_2 are unicycle graphs, but only $\Psi(G_2)$ is a greedoid.

For instance, $S_1 = \{a, b, d\}$ and $S_2 = \{b, c, d\}$ belong to $\Psi(G_1)$, where G_1 is the graph in Figure 2, but only S_1 has an accessibility chain, namely, $\{a\} \subset \{a, d\} \subset S_1$. Nevertheless, having a simplicial vertex is a necessary but not a sufficient condition for a stable set to admit an accessibility chain; e.g., $S_2 = \{b, c, d\} \in \Psi(G_1)$ has a pendant vertex and no accessibility chain. However, there exist graphs where every maximum stable set has an accessibility chain, e.g., the graph G_2 from Figure 2.

Evidently, if $\Psi(G)$ has the accessibility property, then every $S \in \Psi(G)$, $|S| = k \geq 2$, has an accessibility chain.

In this paper we characterize the unicycle graphs whose family of local maximum stable sets are greedoids. Namely, we demonstrate that if C_k is the unique cycle of G , then the family $\Psi(G)$ is a greedoid for $k = 3$, while for $k \geq 4$, $\Psi(G)$ is a greedoid if and only if either (a) k is an even number and all maximum matchings of G are uniquely restricted, or (b) k is an odd number and the closed neighborhood of every local maximum stable set of G induces a König-Egerváry graph.

2 Results

Let C be the unique cycle of a graph G . Clearly, for every $e \in E(C)$, the resulting graph $G - e$ is a forest.

Theorem 2.1 [10] *For any forest T , $\Psi(T)$ is a greedoid on its vertex set.*

This assertion fails for general graphs, and even for unicycle graphs is not always true (e.g., see the graph G in Figure 3, whose family $\Psi(G)$ is not a greedoid).

Proposition 2.2 *If $\Psi(G)$ is a greedoid, then $\Omega(C_k) \cap \Psi(G) = \emptyset$ for its every induced cycle C_k of size $k \geq 4$.*

Proof. Suppose that there exists $S \in \Omega(C_k) \cap \Psi(G)$ in G for some $k \geq 4$. Since $\Psi(G)$ is a greedoid, there is a chain of local maximum stable sets

$$\{x_1\} \subset \{x_1, x_2\} \subset \dots \subset \{x_1, x_2, \dots, x_{k-1}\} \subset \{x_1, x_2, \dots, x_q\} = S,$$

where $q = |S| \geq 2$. Hence, x_1 must be a pendant vertex in G , contradicting the fact that x_1 belongs to $V(C_k)$. ■

The graph G from Figure 3 satisfies the condition that $\Omega(C_k) \cap \Psi(G) = \emptyset$ for its every cycle C_k of size $k \geq 4$. Nevertheless, $\Psi(G)$ is not a greedoid, since $S = \{a, d, g\} \in \Psi(G)$, while S admits no accessibility chain. In other words, the converse of Proposition 2.2 is not true.




Figure 3: $S = \{a, d, g\} \in \Psi(G)$, but S admits no accessibility chain in G .

In the sequel, we distinguish between the following cases: $C = C_3$ and $C = C_k$, $k \geq 4$.

Theorem 2.3 *If G is a graph that has a C_3 as its unique cycle, then $\Psi(G)$ is a greedoid.*

Proof. Let $V(C_3) = \{x_i : 1 \leq i \leq 3\}$.

If $V(G) = V(C_3)$, then it is easy to see that $\Psi(G)$ is a greedoid.

Let $V(G) \neq V(C_3)$. According to Theorem 1.1, it is sufficient to show that $\Psi(G)$ satisfies the accessibility property. In other words, we have to build an accessibility chain for any $S \in \Psi(G)$.

Let $T_{ij}, i \in \{1, 2, 3\}, j \in \{0, 1, \dots, n_i\}$ be subtrees of G such that: $T_{i0} = (\{x_i\}, \emptyset)$, while for $j \geq 1$, T_{ij} is joined by an edge to $x_i, i \in \{1, 2, 3\}$, respectively (whenever such a subtree exists).

- *Case 1.* $S \cap V(C_3) = \emptyset$.

Then $S \in \Psi(G - \{x_1, x_2, x_3\})$ and $T = G - \{x_1, x_2, x_3\}$ is a forest. Therefore, by Theorem 2.1, there is an accessibility chain for S in T . This is an accessibility chain for S in G , as well, because the neighborhoods of the sets belonging to the chain are the same in T and G .

- *Case 2.* $S \cap V(C_3) \neq \emptyset$, e.g. $S \cap V(C_3) = \{x_1\}$.

Let us denote

$$\begin{aligned} G_1 &= G[\cup\{V(T_{1j}) : 0 \leq j \leq n_1\}], & S_1 &= S \cap V(G_1), \\ G_2 &= G[\cup\{V(T_{2j}) : 1 \leq j \leq n_2\}], & G_3 &= G[\cup\{V(T_{3j}) : 1 \leq j \leq n_3\}], \\ S_{ij} &= S \cap V(T_{ij}), 1 \leq j \leq n_i, & S_i &= \cup\{S_{ij}, 1 \leq j \leq n_i\}, i = 2, 3. \end{aligned}$$

Claim 1. S_2, S_3 are local maximum stable sets in G_2, G_3 , respectively.

Otherwise, suppose that, for instance, $S_2 \notin \Psi(G_2)$. Then, there is $W \in \Omega(N_{G_2}[S_2])$ with $|S_2| < |W|$. Since $N_{G_2}[S_2] \subseteq N_G[S_2]$ and $x_1 \notin N_G[S_2]$, it follows that $S_1 \cup W \cup S_3$ is a stable set in $N_G[S]$, but larger than S , in contradiction to the choice $S \in \Psi(G)$.

Claim 2. $S_1 \in \Psi(G_1)$.

Otherwise, there must exist some stable set $W \subseteq N_{G_1}[S_1]$ with $|S_1| < |W|$. Hence, W is also stable in G , and since $N_{G_1}[S_1] \subseteq N_G[S_1]$, it follows that $W \cup S_2 \cup S_3$ is a stable set in $N_G[S]$, but larger than S , in contradiction to the choice $S \in \Psi(G)$.

Claim 3. There is an accessibility chain of S in G .

We distinguish between the following two cases.

Case 3.1. $S_1 - \{x_1\} \in \Psi(G_1)$.

Hence, we infer that $S_1 - \{x_1\} \in \Psi(G_1 - \{x_1\})$, which together with *Claim 1* imply

$$(S_1 - \{x_1\}) \cup S_2 \cup S_3 \in \Psi(G - \{x_1, x_2, x_3\}).$$

Therefore, according to Theorem 2.1, there exists an accessibility chain for the local maximum stable set $(S_1 - \{x_1\}) \cup S_2 \cup S_3$ in $T = G - \{x_1, x_2, x_3\}$, because T is a forest. This is an accessibility chain for $(S_1 - \{x_1\}) \cup S_2 \cup S_3$ in G , as well, because the neighborhoods of the sets belonging to the chain are the same in T and G . Clearly, this gives rise to an accessibility chain for $S_1 \cup S_2 \cup S_3$.

Case 3.2. $S_1 - \{x_1\} \notin \Psi(G_1)$.

Since $S_1 \in \Psi(G_1)$ (by *Claim 2*) and $\Psi(G_1)$ is a greedoid (by Theorem 2.1), there is $v \in S_1$, such that $S_1 - \{v\} \in \Psi(G_1)$.

We assert that $(S_1 - \{v\}) \cup S_2 \cup S_3 \in \Psi(G)$.

Otherwise, there is a stable set $A \subset N_G[(S_1 - \{v\}) \cup S_2 \cup S_3]$ with

$$|A| > |S_1| - 1 + |S_2| + |S_3|.$$

Therefore, either $x_2 \in A$ or $x_3 \in A$. Without lack of generality suppose that $x_2 \in A$. Hence, there exists a stable set $B \subset N_G[S_2]$ in G_2 such that $|B| = |S_2|$ and $B \cup \{x_2\}$ is stable in G . Since $S_1 - \{x_1\} \notin \Psi(G_1)$, there exists $S_0 \subset N[S_1 - \{x_1\}]$, such that $|S_0| > |S_1 - \{x_1\}|$. Clearly, $x_1 \notin S_0 \subset N[S_1]$. Consequently, $S_0 \cup B \cup \{x_2\} \cup S_3$ is a stable set in $N_G[S]$ of size greater than $|S|$, and that contradicts the choice of $S \in \Psi(G)$.

Thus, we obtain

$$S, (S_1 - \{v\}) \cup S_2 \cup S_3 \in \Psi(G),$$

$$x_1 \in S_1 - \{v\} \in \Psi(G_1), \quad S_2 \in \Psi(G_2), \quad S_3 \in \Psi(G_3),$$

and

$$S = S_1 \cup S_2 \cup S_3 \supset (S_1 - \{v\}) \cup S_2 \cup S_3.$$

Now, if $(S_1 - \{v\}) - \{x_1\} \in \Psi(G_1)$, we continue as in *Case 1*, which leads immediately to an accessibility chain.

Otherwise, we find some vertex

$$v' \in S_1 - \{v\} \in \Psi(G_1), v' \neq x_1,$$

such that $(S_1 - \{v\}) - \{v'\}$ belongs to $\Psi(G_1)$, and we can continue, as in *Case 2*, to increase the size of the chain of local maximum stable sets we are building one by one.

Since the set S is finite, at the end of the above procedure we obtain an accessibility chain of S in G . ■

Let us remark that the graph G in Figure 3 is a unicycle bipartite graph, and as we mentioned before, its family $\Psi(G)$ is not a greedoid. However, there exist bipartite graphs whose families $\Psi(G)$ are greedoids.

Trying to characterize these bipartite graphs, we found out an interesting connection between their local maximum stable sets of a graph and their matchings, but of some special kind [9, 10, 11, 12].

A *perfect matching* is a matching saturating all the vertices of the graph. A matching M of a graph G is called a *uniquely restricted matching* if M is the unique perfect matching of the subgraph induced by the vertices it saturates [4].

Recall that a cycle C is *alternating* with respect to a matching M if for any two incident edges of C exactly one of them belongs to M , [7]. It is clear that an M -alternating cycle should be of even size.

Theorem 2.4 [4] *A matching M in a graph G is uniquely restricted if and only if G does not contain an alternating cycle with respect to M .*

Notice that the graph G in Figure 3 has maximum matchings that are not uniquely restricted. It turns out that the existence of such matchings is the real reason why $\Psi(G)$ is not a greedoid.

Theorem 2.5 [11] *For a bipartite graph G , the family $\Psi(G)$ is a greedoid if and only all maximum matchings of G are uniquely restricted.*

According to Theorem 2.4, all maximum matchings of a unicycle non-bipartite graph G are uniquely restricted.

Nevertheless, it is not sufficient for $\Psi(G)$ to be a greedoid. For example, the graphs G_1, G_2 in Figure 4 are unicycle non-bipartite graphs, $\Psi(G_2)$ is a greedoid, while $\Psi(G_1)$ is not a greedoid, because $\{u, v\} \in \Psi(G_1)$, but $\{u\}, \{v\} \notin \Psi(G_1)$.



Figure 4: Non-bipartite triangle-free graphs with unique perfect matchings.

A *triangle-free graph* is a graph having no induced subgraph isomorphic to C_3 . The graphs in Figure 4 are triangle-free König-Egerváry graphs, while $G_1[\{u, v\}]$ is not a König-Egerváry graph. The existence of such a subgraph is the reason for $\Psi(G_1)$ not to be a greedoid.

Theorem 2.6 [12] *If G is a triangle-free graph, then $\Psi(G)$ is a greedoid if and only if all its maximum matchings are uniquely restricted and the closed neighborhood of each local maximum stable set of G induces a König-Egerváry graph.*

Now, using the fact that, by Theorem 2.4, all the maximum matchings of a graph without even cycles must be uniquely restricted, and combining Theorems 2.3, 2.5, 2.6, we conclude with the following.

Theorem 2.7 *Let C_k be the unique cycle of the graph G . Then, the following statements are true:*

- (i) *if $k = 3$, then $\Psi(G)$ is a greedoid;*
- (ii) *if $k = 2q \geq 4$, then $\Psi(G)$ is a greedoid if and only if all maximum matchings of G are uniquely restricted;*
- (iii) *if $k = 2q + 1 \geq 5$, then $\Psi(G)$ is a greedoid if and only if the closed neighborhood of every local maximum stable set of G induces a König-Egerváry graph.*

3 Conclusions

In this paper we have completely characterized the unicycle graphs whose families of local maximum stable sets form greedoids on their vertex sets. In [9] we showed that even unicycle graphs whose families of local maximum stable sets are greedoids can be recognized in polynomial time. The key question that remains open is whether there exists a polynomial time recognition algorithm for odd unicycle graphs whose families of local maximum stable sets are greedoids.

References

- [1] A. Björner, G. M. Ziegler, *Introduction to greedoids*, in N. White (ed.), *Matroid Applications*, 284-357, Cambridge University Press, 1992.
- [2] R. W. Deming, *Independence numbers of graphs - an extension of the König-Egerváry theorem*, Discrete Mathematics **27** (1979) 23-33.
- [3] E. Egerváry, *On combinatorial properties of matrices*, Mat. Lapok **38** (1931) 16-28.
- [4] M. C. Golumbic, T. Hirst, M. Lewenstein, *Uniquely restricted matchings*, Algorithmica **31** (2001) 139-154.
- [5] B. Korte, L. Lovasz, R. Schrader, *Greedoids*, Springer-Verlag, Berlin, 1991.
- [6] D. König, *Graphen und Matrizen*, Mat. Lapok **38** (1931) 116-119.
- [7] S. Krogdahl, *The dependence graph for bases in matroids*, Discrete Mathematics **19** (1977) 47-59.
- [8] V. E. Levit, E. Mandrescu, *On the structure of α -stable graphs*, Discrete Mathematics **236** (2001) 227-243.
- [9] V. E. Levit, E. Mandrescu, *Unicycle bipartite graphs with only uniquely restricted maximum matchings*, in Proceedings of the Third International Conference on Combinatorics, Computability and Logic, (DMTCS'1), Springer, (C.S. Calude, M. J. Dinneen and S. Sburlan eds.) (2001) 151-158.
- [10] V. E. Levit, E. Mandrescu, *A new greedoid: the family of local maximum stable sets of a forest*, Discrete Applied Mathematics **124** (2002) 91-101.
- [11] V. E. Levit, E. Mandrescu, *Local maximum stable sets in bipartite graphs with uniquely restricted maximum matchings*, Discrete Applied Mathematics **132** (2003) 163-174.
- [12] V. E. Levit, E. Mandrescu, *Triangle-free graphs with uniquely restricted maximum matchings and their corresponding greedoids*, Discrete Applied Mathematics **155** (2007) 2414-2425.
- [13] V. E. Levit, E. Mandrescu, *On local maximum stable sets of the corona of a path with complete graphs*, Proceedings of the 6th Congress of Romanian Mathematicians, June 28 - July 4, 2007, University of Bucharest, Romania (in press).

- [14] V. E. Levit, E. Mandrescu, *Well-covered graphs and greedoids*, Proceedings of the 14th Computing: The Australasian Theory Symposium (CATS2008), Wollongong, NSW, Australia, Conferences in Research and Practice in Information Technology Volume **77** (2008) 89-94.
- [15] V. E. Levit, E. Mandrescu, *The clique corona operation and greedoids*, Combinatorial Optimization and Applications, Second International Conference, COCOA 2008, Lecture Notes in Computer Science **5165** (2008) 384-392.
- [16] V. E. Levit, E. Mandrescu, *Graph operations that are good for greedoids*, MOPTA 2008, University of Guelph, Guelph, Canada, arXiv:0809.1806v1[math.CO] (2008) 9 pp.
- [17] V. E. Levit, E. Mandrescu, *Interval greedoids and local maximum stable sets in graphs*, Los Alamos Archive, arXiv:0811.4089v1[math.CO] (2008) 13 pp.
- [18] G. L. Nemhauser, L. E. Trotter, Jr., *Vertex packings: structural properties and algorithms*, Mathematical Programming **8** (1975) 232-248.
- [19] F. Sterboul, *A characterization of the graphs in which the transversal number equals the matching number*, Journal of Combinatorial Theory Series B **27** (1979) 228-229.